



H-WV CP Violation Measurement Update

Ian Anderson¹, Sara Bolognesi¹, Fabrizio Caola¹, **Yanyan Gao²**,
Andrei Gritsan¹, Kirill Melnikov¹,
Markus Schulze³, Nhan Tran², Andrew Whitbeck¹ and Yaofu Zhou¹

¹ Johns Hopkins University, ² Fermilab, ³ Argonne National Lab


Snowmass Meeting in Higgs group, Minneapolis, August 2nd, 2013

Update in e+e- Collider Experiments

- Compare to last update from A.Whitbeck in Seattle <http://goo.gl/XZXdL>
 - Extend the studies for 350 GeV and 1 TeV
 - Use MC simulated events in the pseudo-experiments
- Significant improvement in e+e- collider experiments compared to LHC
 - Presented at EPS by Yanyan Gao: <http://goo.gl/oalvdQ>
 - In the LHC analysis, explore only the decay kinematics of $X \rightarrow ZZ \rightarrow 4l$


CP violation $\sim 3\sigma$ discovery potential

	\sqrt{s} (Lumi)	f_3	f_{a3}	$\sigma(f_{a3})$
LHC $X \rightarrow ZZ(4l)$	14TeV(300/fb)	0.18	0.18	0.06
	14TeV(3000/fb)	0.06	0.06	0.02
$e^+e^- \rightarrow Z^* \rightarrow Z(l)X(bb)$	250GeV(250/fb)	0.1	2.10E-03	7.00E-04
	350GeV(350/fb)	0.1	3.36E-04	1.12E-04
	500GeV(500/fb)	0.1	1.05E-04	3.90E-05
	1TeV(1000/fb)	0.1	1.95E-05	7.22E-06



Development on the $\text{pp} \rightarrow \text{H}$ in WBF Channel

- Significant boost in relative 0-/0+ cross-section due to the off-shell-ness of the W/Z-boson



$$D_{0-}^{WBF} = \left[1 + \frac{P_{0-}^{WBF}}{P_{0+}^{WBF}} \right]^{-1}$$

- ME based on VBF jets and Higgs 4-momenta
 - Method can be used for any VBF production
 - Preliminary study uses the VBF $H \rightarrow ZZ \rightarrow 4l$

CP violation $\sim 3\sigma$ discovery potential

	\sqrt{s} (Lumi)	f_3	f_{a3}	$\sigma(f_{a3})$
LHC $X \rightarrow ZZ(4l)$	300/fb	0.18	0.18	0.06
	3000/fb	0.06	0.06	0.02
LHC $VBF \rightarrow X \rightarrow ZZ(4l)$	300/fb (jet pt > 30)	0.5	0.0084	0.003
	3000/fb (jet pt > 30)	0.15	0.0015	0.0005
	3000/fb (jet pt > 50)	0.5	0.0049	0.0015

